




# My Asthma Plan



Your asthma plan tells you when to take your asthma medicines.

And what to do when your asthma gets worse.



Name: \_\_\_\_\_

## 1. My daily asthma medicines

- My preventer inhaler is called \_\_\_\_\_ and its colour is \_\_\_\_\_
- I take \_\_\_\_\_ puff/s of my preventer inhaler in the morning and \_\_\_\_\_ puff/s at night. I do this every day even if I feel well.
- Other asthma medicines I take every day:  
\_\_\_\_\_  
\_\_\_\_\_
- My reliever inhaler is called \_\_\_\_\_ and its colour is \_\_\_\_\_. I take \_\_\_\_\_ puff/s of my reliever inhaler (usually blue) when I wheeze or cough, my chest hurts or it's hard to breathe.
- My best peak flow is \_\_\_\_\_

## 2. When my asthma gets worse

I'll know my asthma is getting worse if:

- I wheeze or cough, my chest hurts or it's hard to breathe, or
- I'm waking up at night because of my asthma, or
- I'm taking my reliever inhaler (usually blue) more than three times a week, or
- My peak flow is less than \_\_\_\_\_

If my asthma gets worse, I should:

Keep taking my preventer medicines as normal.

I also take \_\_\_\_\_ puff/s of my reliever inhaler (usually blue) every four hours.

If I'm not getting any better doing this I should see my doctor or asthma nurse today.

Does doing sport make it hard to breathe?




If YES

I take:

\_\_\_\_\_ puff/s of my reliever inhaler (usually blue) beforehand.

Remember to use my inhaler with a spacer (if I have one)





# My Asthma Plan

## 3. When I have an asthma attack

### I'm having an asthma attack if:

- My reliever inhaler (usually blue) isn't helping, or
- I can't talk or walk easily, or
- I'm breathing hard and fast, or
- I'm coughing or wheezing a lot, or
- My peak flow is less than \_\_\_\_\_

### When I have an asthma attack, I should:

Sit up – don't lie down. Try to be calm.

Take one puff of my reliever inhaler every 30 to 60 seconds up to a total of 10 puffs.

Even if I start to feel better, I don't want this to happen again, so I need to see my doctor or asthma nurse today.



If I still don't feel better and I've taken ten puffs, I need to call 999 straight away. If I am waiting longer than 15 minutes for an ambulance I should take another \_\_\_\_\_ puff/s of my reliever inhaler (usually blue) every 30 to 60 seconds (up to 10 puffs).



You and your parents can get your questions answered:

Call Asthma UK's friendly Helpline

Monday to Friday 9am to 5pm

**0300 222 5800**

Get information at

**[www.asthma.org.uk](http://www.asthma.org.uk)**

### My asthma triggers (things that make my asthma worse)

---



---



---



---

Make sure you have your reliever inhaler (usually blue) with you. You might need it if you come into contact with things that make your asthma worse.

### I need to see my asthma nurse every six months

Date I got my asthma plan:

---

Date of my next asthma review:

---

Doctor/asthma nurse contact details:

---

---

---

### Parents – get the most from your child's action plan

Make it easy for you and your family to find it when you need it

- Take a photo and keep it on your mobile (and your child's mobile if they have one)
- Stick a copy on your fridge door
- Share your child's action plan with school, grandparents and babysitter (a printout or a photo).